

Analysis of a Fossil Bone from the Archaeological Settlement Malu Rosu, Romania by Accelerator Mass Spectrometry

Agata Olariu (agata@ifin.nipne.ro)

*National Institute for Physics and Nuclear Engineering, PO Box MG-6, 76900
Magurele, Bucharest, Romania*

Ragnar Hellborg (ragnar.hellborg@nuclear.lu.se) and Kristina
Stenström (kristina.stenstrom@nuclear.lu.se)

*Department of Nuclear Physics, Lund University, Sölvegatan 14, SE-223 62
Lund, Sweden*

Göran Skog (goran.skog@c14lab.lu.se)

Department of Quaternary Geology, Lund, Tornavägen 13, SE-223 63 Lund

Mikko Faarinen (mikko.faarinen@nuclear.lu.se), Per Persson
(per.persson@nuclear.lu.se) and Bengt Erlandsson
(bengt.erlandsson@nuclear.lu.se)

Department of Nuclear Physics, Lund University

Ion V. Popescu (ivpopes@ifin.nipne.ro)

National Institute for Physics and Nuclear Engineering, Bucharest

Emilian Alexandrescu

Institute of Archaeology, Bucharest



Abstract. A fossil bone from the archaeological site Malu Rosu Giurgiu, in Romania has been analyzed by accelerator mass spectrometry to estimate its age by determining its ^{14}C content. The radiocarbon age of the bone is in agreement with the age obtained by the method for age determination, based on fluorine content. This is the first radiocarbon dating for the final Neolithic period, for this archaeological settlement in the Romanian region.

1. Introduction

Among the physical methods of importance for archaeological applications, dating techniques occupy a special place. Historians need a variety of information regarding their objects: structure, provenance, culture or authenticity^{1,2}. At the same time there is a great need of dating with precise and objective techniques of the historically valuable material. There are two categories of physical methods of dating, namely one in which the measured quantity decreases with time, and one in which the physical quantity accumulates in the sample with time. To the first category belong the methods in which the information on age is given by the disintegration rate of a radioactive nucleus, ranging from ^{40}K , Th and U (relatively abundant isotopes) to more rare isotopes such as ^{14}C , ^{10}Be , ^{26}Al , ^{32}Si , ^{36}Cl , ^{41}Ca , ^{53}Mn , ^{210}Pb . The second category includes the methods that are based on the measurement of the accumulated defects produced by radiation in the

environment, using various techniques: thermoluminescence, electron paramagnetic resonance, fission traces in rocks.

Among dating methods, radiocarbon dating is the most frequently used. In recent decades accelerator mass spectrometry (AMS)^{3,4}, which constitutes a highly sensitive method for counting atoms, has been used for the detection of ^{14}C . The fact that AMS counts atoms and not decays results in great advantages compared to radiometric techniques, requiring smaller samples and shorter measuring times.

In the present work we have studied using the AMS technique a fossil bone found in the course of the archaeological excavation of a workshop for the processing of flintstones at Malu Rosu, Giurgiu, near the Danube River, in Romania. This archaeological assembly is believed to have an age of about 20,000 years, as based on an historical assessment. Other specimens of fossil bones from Malu Rosu have earlier been analyzed by a relative chemical method. The age was estimated by determining the fluorine content⁵ using neutron activation analysis (NAA).

2. Experimental

2.1. PREPARATION OF THE BONE SAMPLE

Pretreatment of the bone sample

The first step concerns the extraction of 'collagen' from the bone structure. Collagen is the native, biochemically intact triple-helical macromolecule. We will use 'collagen' to refer to collagen that has undergone a degree of diagenesis. Alterations during diagenesis are considered to include humification of parts of the molecule, attachment of exogenous humic materials, and hydrolysis with loss of some amino acids. We have essentially applied the Longin method⁷:

A dried bone specimen of about 1 g is powdered using a hard knife. The bone powder is put into a glass vessel with a rubber cork, which is then connected to a vacuum pump. About 70 - 80 ml of 2% HCl is added in drops to the bone powder to remove the carbonates. The CO₂ released is evacuated by pumping for 2 - 5 minutes. The bone is left in this solution for 15 - 24 h. 'Collagen' is then retrieved as a sediment in the vessel. The acid is poured off and the bone sample is washed with distilled water three times (until the solution is neutral).

The 'collagen' is transferred to a 50 ml vessel. 20 - 30 ml of distilled water is added together with some drops of 2% HCL to obtain a pH in the range 2 - 3. This solution is kept at a temperature of 90 °C overnight. The pH must remain in the range 2 - 3 for the whole period of heating, which ensures that the 'collagen' is dissolved.

The 'collagen' solution is filtered to remove all the impurities, and put in the oven at 110 °C until is completely evaporated. The 'collagen', in gelatin form is thus concentrated at the bottom of the test tube.

Transformation to elemental carbon

The next step in the preparation of the bone sample is the transformation of the 'collagen' into pure carbon in the experimental set-up, presented in detail in refs. 6, 8 and 9. The conversion to elemental carbon is done in two steps: the formation of CO₂ by the combustion of the 'collagen', and the reduction of CO₂ to carbon.

Combustion. About 200 mg of CuO as oxidation agent is added to the 'collagen' sample in the test tube in order to attain a 100% combustion. The system is evacuated and the test tube containing the sample is heated with a flame. The water vapor that is released during the combustion is caught in a cold trap.

Reduction. About 10 mg of metallic iron is put into a double-legged test tube as catalyst for the reduction of the CO₂ to carbon. Prior to the reduction process, the catalyst is heated under vacuum to 650 °C. With CO₂ transferred to and frozen in the cold trap of the reduction part of the apparatus, hydrogen gas is let into the system. It is suitable to have 3 times more hydrogen gas than carbon dioxide. The reduction

process takes about 4 h. When the reduction is complete, the carbon sample is dried by pumping for about 20 minutes.

The carbon sample and the iron catalyst are pressed into a copper holder with a stainless steel piston. The iron catalyst provides a good thermal conduction of the sample, which is important for the sputtering process into the ion source.

The copper holder with the pure carbon sample is put into the ion source of the accelerator.

2.2. THE AMS ANALYSIS AT THE TANDEM ACCELERATOR

The sample of bone, as pure carbon in the copper holder is arranged in a wheel with 20 positions in the following sequence: 2 standards of oxalic acid, 1 standard of anthracite, 15 carbon samples, with the other 2 positions being for a graphite and an optimizing sample. Anthracite, being a very old coal is considered to contain no traces of ^{14}C traces and by its measurement the background for ^{14}C is obtained both of the accelerator and of the preparation procedure of the samples. Oxalic acid is a standard SRM prepared by the USA National Bureau of Standards, with an accurately known activity of ^{14}C . The oxalic acid is used to normalize the value of the $^{14}\text{C}/^{13}\text{C}$ ratio of the sample. The wheel with samples and standards is put into the ion source of the accelerator. The central part of the Lund AMS system is a Pelletron tandem accelerator, model 3UDH, produced by NEC, Wisconsin USA, and is shown in Fig. 1. The accelerator is run at 2.4 MV during AMS experiments, which is optimal for the C^{3+} charge state when using $3\text{ }\mu\text{g}/\text{cm}^2$ carbon stripper foils. On the experimental beam line

a magnetic quadrupole triplet, a velocity filter and a second analyzing magnet have been installed. The $^{13}\text{C}^{3+}$ beam current is usually ≈ 80 nA.

The particle identification and measuring system consists of a silicon surface barrier detector of $\Phi=25$ mm. Part of the accelerator system is operated and controlled by a PC-based system. The computer system analyses alternately the data of the ^{13}C current received from a current integrator and the ^{14}C counts arriving from the particle detector, to obtain, finally, the ratio $^{14}\text{C}/^{13}\text{C}$ for each sample. This ratio is corrected for the background (obtained from the anthracite and oxalic acid standards) for every sample. Each sample has been measured 7 times. The precision of the measurements is around 1 % .

3. Results and Discussions

3.1. THE RADIOCARBON AGE

The age of the bone was determined in two steps: first the *conventional radiocarbon age* was calculated and then the age was converted to calendar years, taking into consideration the fluctuations of the ^{14}C concentration in the atmosphere over time.

The conventional radiocarbon age was calculated by the equation ^{10,11,12}:

$$\text{Age} = -8033 \ln \left(\frac{(^{14}\text{C}/^{13}\text{C})_{S[-25]}}{0.9558 (^{14}\text{C}/^{13}\text{C})_{Ox[-19]}} \right) \text{ years BP} \quad (1)$$

where $(^{14}\text{C}/^{13}\text{C})_{S[-25]}$ is the isotopic ratio for the sample, corrected for the background, normalized to $\delta^{13}\text{C}=-25$ ‰ and $(^{14}\text{C}/^{13}\text{C})_{Ox[-19]}$ is the isotopic ratio for the oxalic acid standard,

corrected for the background, normalized to $\delta^{13}C = -19\text{‰}$,
with $\delta^{13}C$ being:


$$\delta^{13}C = \left[\frac{R}{R_{PDB}} - 1 \right] \times 1000\text{‰} \quad (2)$$

where R is the $^{13}C/^{12}C$ isotopic ratio for the sample and the oxalic acid, respectively, and R_{PDB} is the $^{13}C/^{12}C$ isotopic ratio of the *PDB* standard. The conventional radiocarbon age of the bone is expressed in years BP, where present is defined to be AD 1950. The conventional radiocarbon age for the analyzed bone from Malu Rosu has been determined to be: 4790 ± 90 BP. The conversion into calendar years was done using the OxCal software¹³ (see Fig. 2), using data from ref. 14. From the processing of the data one obtained the age of the bone from Malu Rosu to be in the interval:

3760 - 3360 BC or 5710 - 5310 BP, with a confidence level of 95.4 % .

3.2. COMPARISON OF THE RADIOCARBON AGE WITH THE AGE ESTIMATED BY FLUORINE CONTENT

The method of dating by measuring the fluorine content has been used since the end of the 19th century. The increase of the fluorine content is the result of the contact of the fossil bone with underground water which contains fluorine^{15,16}. The main inorganic component of the bone, hydroxyapatite is slowly altered by the exchange of hydroxyl ions (OH^-) with ions F^- :



The result of this exchange is the formation of fluoroapatite $\text{Ca}_{10}(\text{PO}_4)_6\text{F}_2$, a compound much more stable than hydroxiapatite.

The rate of accumulation of fluorine in bones depends on the environmental conditions of the burials, the temperature and the content of fluorine in underground waters.

Some specimens of bones from Malu Rosu have been analyzed before by NAA to determine their fluorine content⁵. The irradiations were done at the VVR-S nuclear reactor at IFIN Bucharest, at a flux of 10^{12} neutrons/cm²·s⁻¹. Samples of bones together with standards of PbF₂ and CaF₂ were irradiated for a period of 4 s, for producing the isotope ²⁰F, with a half life of $T_{1/2}=11$ s and the main γ -ray of 1633.6 keV. The periods of decays and times of measurement for the sample were: $t_{decay}=40$ s, $t_{meas}=20$ s, and for the standard: $t_{decay}=80$ s, $t_{meas}=20$ s. For 8 specimens of bones from Romanian regions, C1, C2, C4, C5, C6, C8 and C9, dated by radiocarbon at Groeningen, the following correlation was established: fluorine concentration - age (radiocarbon). For other bones: B1 (cave bear), B2 (cave bear), B3 (cave bear), B4 (mammoth) and MR (animal bone from Malu Rosu) was evaluated the age, from the content of fluorine, determined by NAA, using the calibration curve. In Fig. 3 are presented the results of NAA of analyzed bones. Also represented are the fluorine concentrations and the ages for some bones from Carribean zone¹³. In this region the climate is warmer and therefore the rate of accumulation of fluorine in the structure of bones is higher than in bones with the same age, from Romanian regions. From the diagram shown in Fig. 3 one can estimate for the bone from Malu Rosu, sample MR, with a content of fluorine of 400 ppm, an age

$\approx 6000 \pm 2000$ years BP. For the bone B1 one could estimate an age around 11000 ± 2000 years BP. For the bear B2 and the mammoth B4, and for the bear B3, one could estimate an inferior limit of 35000, respectively 37000 years BP, given by the calibration curve.

Taking into consideration a possible saturation of the curve, the bones B2 and B4, and B3 could have an age of around 40000, respectively 45000 years BP.

The age of the bone from Malu Rosu of 5510 ± 200 BP, determined by ^{14}C measurements using AMS in the present study is in agreement with the estimated value of the age of 5000 ± 1000 years BP, determined in other fossil bones from the Malu Rosu, by measuring their content of fluorine, using NAA.

3.3. ARCHAEOLOGICAL CONSIDERATIONS

The settlement at Malu Rosu Giurgiu is situated in the south of Romania, on the bank of the River Danube, in a region where the inferior terrace is preserved intact.

The stratigraphy of the settlement is very complex, containing, besides the four cultural Aurignacian levels (workshops for processing flintstones have been found with some hearths in the central parts) and cultural levels from the final middle Neolithic (Boian culture), Eneolithic (Gumelnița culture), la Tène period, early Middle Age (Dridu culture, IX-XI centuries) (Fig. 4).

Due to the specific conditions, namely open areas, the vegetal deposit and the loessial deposit, which start at a depth of approximately one metre, are affected by rodent passages both contemporary and fossil, reaching in some cases a depth of 6 m. Under these conditions some archaeological objects have been disturbed and moved from their

initial position. Furthermore, the Neolithic, Eneolithic and Medieval archaeological complexes have sometimes affected the Aurignacian cultural layers.

The fragment of bone studied by AMS in the present study originates from the superior part of an Aurignacian layer I ¹⁷, Surface III/1995, the workshop for the processing of flintstones Nr. 1, square D/11, depth 1.98 m (Fig. 4). The inferior part of this Aurignacian layer, the oldest one in Malu Rosu, has been dated by AMS, at Groeningen, on the basis of charcoal fragments from hearths, to have an age of $21,140 \pm 120$ BP (GrA-5094) and $22,790 \pm 130$ BP (GrA-6037).

The value of 3560 ± 200 BC for the age of the bone determined by AMS in the present work associates the bone with the Neolithic level of the Malu Rosu settlement, attributed by us to the final stage of evolution of the Boian culture. In the layer mentioned one finds, at a depth of between 0.80 - 1.10 m, the archaeological remains of three houses belonging to the final stage of the Boian culture.

Some radiocarbon datings for the Boian culture can be quoted for some settlements from the Danubian region: for the Cascioarele settlement, with a layer dated at 4035 ± 125 BC (BIN-800) and 3620 ± 100 BC (BIN-796) and the settlement at Radovanu, from the oldest layer (transitional stage to Eneolithic), dated to 3820 ± 100 (BIN-1233)¹⁸.

Thus the fossil bone analyzed in the present work, found in the Palaeolithic layer of the Malu Rosu settlement is dated to have an age of 3560 ± 200 BC, and belongs in fact to the Neolithic layer, situated above the Palaeolithic one. This result can be explained by the exis-

tence of a dynamics of archaeological objects, by which they can be moved from their original places by an interference with the biosphere.

By dating the bone from Malu Rosu, Giurgiu one obtains the first radiocarbon estimation of age for the final Neolithic period, for this archaeological settlement in the Romanian region.

References

1. J. W. Michels, Dating Methods in Archaeology, Seminar Press, 1973
2. G. A. Wagner, Archaeometric Dating, in Lectures in Isotope Geology,
eds. E. Jager and J. C. Hunziker, Springer, 1979, p. 178
3. W. Kutschera, *Accelerator mass spectrometry: counting atoms rather than decays*, Nuclear Physics News, Vol. **3**, No. **1** (1993) 15
4. C. Tuniz, J. Bird, D. Fink, G. F. Herzog, Accelerator Mass Spectrometry, Ultrasensitive Analysis for Global Science, CRC Press, 1998
5. C. Besliu and A. Olariu, I. Popescu and T. Badica, *Microelements in fossil bones and the estimation of age*, SPIE, **2339** (1995) 487, and 4th International Conference on Applications of Nuclear Techniques: Neutrons and their Applications., Crete, Greece, 12-18 June, 1994
6. A. Olariu, R. Hellborg, K. Stenström, G. Skog, M. Faarinen, P. Persson, B. Erlandsson, I. V. Popescu, E. Alexandrescu, *Analysis of a Fossil Bone from the Archaeological Settlement Malu Rosu, Romania by accelerator mass spectrometry*, Report 08/00, LUNDFD6/(NFFR-3081)/1-30/(2000), Lund, 2000
7. R. Longin, *New method of collagen extraction for radiocarbon dating*, Nature, v **230** (1990) p. 241-242
8. J. S. Vogel, J. R. Southon, D. E. Nelson and T. A. Brown, Nucl. Instr. and Meth. **B5** (1984) p. 289

9. K. Stenström, G. Skog, B. Erlandsson, R. Hellborg, A. Jänis, A. Wiebert, *A sample preparation system for production of elemental carbon for AMS analyses*, Report 02/94, LundFD6/(NFFR-3065)/1-33/(1994), Lund 1994

10. R. Gillespie, Radiocarbon User's Handbook, Oxford University, Committee for Archaeology, Monograph Number Three, Oxonian Rewley Press, 1984
11. D. J. Donahue, T. W. Linick and A. J. T. Jull, *Isotope ratio and background corrections for accelerator mass spectrometry radiocarbon measurements*, Radiocarbon, Vol. **32**, No. **2** (1990) p. 135
12. M. Stuiver, H. Polach, *Reporting of ^{14}C Data*, Radiocarbon, Vol. **19**, No. **3** (1977) p. 333-363
13. C. Ramsey, from Oxford Radiocarbon Accelerator Unit, UK, OxCal software, http://www.rlaha.ox.ac.uk/oxcal/oxcal_h.html
14. M. Stuiver, Radiocarbon, Vol. **40**, No. **3** (1988)
15. Z. Goffer, Archaeological Chemistry, John Wiley And Sons. Inc. N.Y., 1980
16. R. P. Parker and H. Toots, *Minor elements in fossil bones*, Geological Society of American Bulletin, Vol. **81** (1970) 925
17. E. Alexandrescu, *A hypothesis about the evolution of the Aurignacian cultural complex from Romanian plane* in Time of History I, Memory and Patrimonium, Bucharest University, 1997, p. 16
18. E. Comsa, The Neolithic on the Romanian territory - considera-

tions,

Publishing House of Academy, Bucharest, Romania, 1987, p. 45.

Figure captions

- Fig. 1. A schematic drawing of the Pelletron system at Lund University, for AMS studies
- Fig. 2. The output diagram produced by the OxCal software for the analyzed bone, showing the conversion of the conventional radiocarbon age to calendar years
- Fig. 3. The diagram of the fluorine concentration versus the radiocarbon age for some fossil bones
- Fig. 4. The archaeological settlement Malu Rosu Giurgiu, Romania: a schematic drawing of the profile of the west wall of SIII, 1995